

## Case Report

### Atypical Presentation of Raynaud's Disease with Inferior Wall Infarct

**Maher M Nasser<sup>1</sup>, Muhammad Fareed Khalid<sup>2</sup>, Mahak Khadija Bhatti<sup>3</sup>**

<sup>1</sup> Clinical Professor, Baylor College of Medicine and Texas Heart Institute, Houston, Texas.

<sup>2,3</sup> Medical Officer, Mayo Hospital, Lahore, Pakistan.

**\*Corresponding author:** Muhammad Fareed Khalid, Mayo Hospital, Lahore, Pakistan.

**Citation:** Muhammad Fareed Khalid (2025) Atypical Presentation of Raynaud's Disease with Inferior Wall Infarct 1(2).

**Copyright:** ©2025 Muhammad Fareed Khalid, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Received:** October 18 2025 | **Accepted:** October 28, 2025 | **Published:** November 08, 2025

**Keywords:** Raynaud's phenomenon, myocardial infarction, vasospastic angina

## Abstract


Raynaud's disease usually presents as blanching of fingers and toes. However, it can present with atypical symptoms too. Vasospastic angina has been associated with Raynaud's disease and can result in anginal episodes. Here, we present a case of 68-year-old patient who presented with an inferior wall infarct and was also exhibiting signs of Raynaud's disease. We also reviewed the available literature about the pathophysiology of these vasospastic changes in patients with Raynaud's disease.

## Introduction

Raynaud Phenomenon (RP) is an exaggerated and episodic vasospasm in response to various stressors such as cold temperature or emotional stress. It presents clinically as sharply demarcated color changes of the skin of the digits and carries a significant burden of pain and hand-related disability. The correlation between Vasospastic Angina and Raynaud's has been explained in several previous observations in the literature. We present a case of an atypical case of Raynaud's with an inferior wall infarct.

## Case Report

Mr. A, a 68-year-old physician, presented with a classic anginal episode manifested as central chest pain radiating to both arms for the past 1 hour. It was associated with shortness of breath and diaphoresis. EKG showed ST segment elevation in inferior leads (Figure 1). The universal diagnostic criterion of STEMI was fulfilled (1). His pain partially resolved after administering sublingual nitroglycerin(2), and subsequent EKG showed no ST elevation.



**Figure 1:** EKG of the patient showing ST segment elevation in leads II, III and AVF.

Over the past ten years, the patient's extremities would show bluish discoloration and blanching when exposed to cold temperatures and while working at his office (Figure 2,3). He also has been having short episodes of chest tightness of moderate severity with no radiation to the arms that were relieved by self-induced cough.



**Figure 2:** Image shows the discoloration of the patient's fingers in response to cold.

**Journal of Clinical Cardiology and Cardiovascular Diagnosis (JCCCD)**

**Figure 3:** Image shows the blanching of patient's fingers when exposed to colder temperatures.

On examination and during one of the episodes, the patient exhibited typical signs of Raynaud's phenomenon, including discoloration and blanching of fingers and toes. However, tibial pulses were intact, and the doppler ultrasound of the lower extremities was normal. His ANA levels showed a titer of 1:160. All other labs including troponin level, electrolytes and renal function tests were normal. Nailfold capillaroscopy was not performed at that time. His cardiovascular risk factors assessment showed no history of smoking, hypertension, hyperlipidemia, or diabetes. However, he had a strong family history of premature coronary artery disease; his father died from an acute myocardial infarction at age 61, and his three brothers died of coronary artery disease in their fifties. The patient had a remote history of *H. pylori* infection.

When a coronary angiogram was done, it revealed 80% occlusion of the distal Right Coronary Artery (RCA) and a 50% lesion in the mid-Left Anterior Descending artery. The RCA was stented, and patient was placed on dual anti-platelets therapy, high-intensity rosuvastatin, low-dose metoprolol, and amlodipine. On follow-up visit one month later, the patient's condition had improved and he was no longer having any episodes of blanching of his finger and toes.

### Discussion

The prevalence of Raynaud's phenomena varies widely between populations. In the US, it is estimated that up to 200,000 people are affected. A higher prevalence among other populations, such as France, is found(3). Raynaud's phenomenon can be either a primary phenomenon or secondary to various underlying medical disorders and drug-related conditions.

Vasospastic angina, previously called Prinzmetal or variant angina, is a form of abnormal vascular response to various stimuli. Prinzmetal et al(4) initially described a clinical syndrome that manifested as rest angina associated with ST-segment elevation that

promptly responded to sublingual nitrates. Since this differs from classical angina described by Heberden (effort angina associated with ST depression (5), he referred to it as variant angina.

The correlation between vasospastic angina and Raynaud's has been explained previously in literature(6,7). Miller et al(6) reported a high prevalence of Raynaud's phenomenon in patients with vasospastic angina occurring at a rate of 24% in patients with variant angina as compared to 3% and 5% in controls (p-value<0.01). Hence vasospastic angina in some patients is a manifestation of a generalized vasospastic disorder and could be associated with migraine, Raynaud's phenomenon, and aspirin-induced asthma(8).

Coronary vasospasm can occur in patients with normal coronary arteries as well as in patients with atherosclerosis. However, it is still poorly understood whether coronary vasospasm can promote atherosclerotic alterations or whether the latter may predispose to spasm(9).

The patient described in our case has a typical history of Raynaud's phenomenon diagnosed clinically, along with recurrent attacks of chest pain relieved by nitrates that were thought to be due to vasospastic angina. However, coronary angiography revealed atherosclerotic narrowing with 80% occlusion of the right coronary artery. This finding was similar to an autopsy result in one of the patients described in Prinzmetal et al paper(4), which revealed that the right coronary artery (which presumably had caused resting angina with ST elevation in inferior leads) had an 80% stenosis. This observation led to the conclusion that coronary spasm was associated with atherosclerosis. Later it was found that coronary spasms can also occur in patients with angiographically normal coronary arteries(10).

Recent experimental studies demonstrate that several endothelium-derived factors mediate vascular smooth muscle cell tone and that damaged endothelium may play a major role in the pathogenesis of vasospasm(11). Endothelial damage may occur even in the early stages of atherosclerosis. Zeiher et al (12) indicated that abnormal vasoconstriction was observed in early atherosclerotic sites in patients without variant angina using intra-coronary ultrasound. Yamagishi et al(13) reported that early stages of atherosclerosis were observed by intracoronary ultrasound at the site of spasm in patients with variant angina, even in the absence of an angiographically significant coronary disease. However, these results were challenged by a porcine model of spasm, indicating that endothelial vasodilator function was preserved at the spastic site(14); it remains to be elucidated.

Journal of Clinical Cardiology and Cardiovascular Diagnosis (JCCCD)

It was suggested that endothelial dysfunction with decreased nitric oxide release and reduced bioavailability combined with vascular smooth muscle cell hyperreactivity might be important in developing coronary spasm. However, endothelial dysfunction alone might not be sufficient to explain vasospastic angina(15). Therefore, more studies need to be done to understand the pathophysiology associated with the vasospasm associated with Raynaud's disease.

**Conclusion:**

This case highlights the importance of taking an extensive history of patient with myocardial infarction so that the root cause can be found. Raynaud's phenomenon can present with an infarct due to atherosclerotic and vasospastic changes. This requires additional treatment of Raynaud's disease along with MI so that further episodes can be prevented.


**References**

- Hegazy MA, Kamal S Mansour KS, Alzyat AS, Mohammad MA, Hegazy AA. (2022). Myocardial Infarction: Risk Factors, Pathophysiology, Classification, Assessment and Management. *Cardiology Research and Reports*. 4(5).
- Bosson N, Isakson B, Morgan JA, Kaji AH, Uner A et al (2019) Safety and Effectiveness of Field Nitroglycerin in Patients with Suspected ST Elevation Myocardial Infarction. *Prehosp Emerg Care*. Sep-Oct;23(5):603-611.
- Maricq HR, Carpentier PH, Weinrich MC, Keil JE, Palesch Y et al (1997) Geographic variation in the prevalence of Raynaud's phenomenon: a 5 region comparison. *J Rheumatol*. May;24(5):879-89. PMID: 9150076.
- Prinzmetal M, Kennamer R, Merliss R, Wada T, BorN.(1959) Angina pectoris. I. A variant form of angina pectoris; preliminary report. *Am J Med*. Sep;27:375-388.
- Silverman ME. William Heberden and Some Account of a Disorder of the Breast. *Clin Cardiol*. 1987 Mar;10(3):211-213
- Miller D, Waters DD, Warnica W, Szlachcic J, Kreeft J (1981) Is variant angina the coronary manifestation of a generalized vasospastic disorder? *N Engl J Med*. Mar 26;304(13):763-766.
- Robertson D, Oates JA.(1978) Variant angina and Raynaud's phenomenon. *Lancet*. Feb 25;1(8061):452.
- Contreras Zuniga E, Gomez Mesa JE, Zuluaga Martinez SX, Ocampo V, Andres UrreaC.(2009) Prinzmetal's angina. *Arq Bras Cardiol*. Aug;93(2):e30-2.
- Ong P, Aziz A, Hansen HS, Prescott E, Athanasiadis A (2015) Structural and Functional Coronary Artery Abnormalities in Patients With Vasospastic Angina Pectoris. *Circ J*;79(7):1431-1438. Epub 2015 Jun 18.
- Cheng TO, Bashour T, Kelsner GA Jr, Weiss L, BacosJ.(1973) Variant angina of Prinzmetal with normal coronary arteriograms. A variant of the variant. *Circulation*. Mar;47(3):476-485.
- Ganz P, Alexander RW. (1985) New insights into the cellular mechanisms of vasospasm. *Am J Cardiol*. Sep 18;56(9):11E-15E.
- Zeiber AM, Schächlinger V, Hohnloser SH, Saurbier B, Just H. (1994) Coronary atherosclerotic wall thickening and vascular reactivity in humans. Elevated high-density lipoprotein levels ameliorate abnormal vasoconstriction in early atherosclerosis. *Circulation*. Jun;89(6):2525-2532.
- Yamagishi M, Miyatake K, Tamai J, Nakatani S, Koyama J et al (1994) Intravascular ultrasound detection of atherosclerosis at the site of focal vasospasm in angiographically normal or minimally narrowed coronary segments. *J Am Coll Cardiol*. Feb;23(2):352-357.
- Miyata K, Shimokawa H, Yamawaki T, Kunihiro I, Zhou X et al (1999) Endothelial vasodilator function is preserved at the spastic/inflammatory coronary lesions in pigs. *Circulation*. Sep 28;100(13):1432-1437.
- Picard F, Sayah N, Spagnoli V, Adjedj J, Varenne O. (2019) Vasospastic angina: A literature review of current evidence. *Arch Cardiovasc Dis*. Jan;112(1):44-55.

**Disclosures**

No relevant conflicts of interest to declare.

[Journal of Clinical Cardiology and Cardiovascular Diagnosis \(JCCCD\)](#)



This work is licensed under Creative Commons Attribution 4.0 License  
DOI:10/JCCCD/2025/09

**Your next submission with OlitesPublishers will reach you the below assets**

- We follow principles of publication led by the Committee on Publication Ethics (COPE).
- Double blinded peer review process which is just as well as constructive.
- Permanent archiving of your article on our website
- Quality Editorial service
- Manuscript accessibility in different formats (PDF, Full Text)
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

Learn more: <https://olitespublishing.com/journal-of-clinical-cardiology-and-cardiovascular-diagnosis/>