

Short Review

Beyond Monoamines: An Integrated Perspective on Antidepressant Activity, Mechanisms, and Emerging Therapeutic Strategies

Samad S, Valencia E, Munguía K, Walker K, Lankin H, Gassman D

Institute of Neuroscience, Mexico

***Corresponding Author:** Munguía K, Institute of Neuroscience, Mexico

Citation: Munguía K, Beyond Monoamines: An Integrated Perspective on Antidepressant Activity, Mechanisms, and Emerging Therapeutic Strategies V2 (2),2026

Copyright: ©2026 Munguía K, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Received date: July 10, 2026; **Accepted date:** July 25, 2026; **Published date:** July 30, 2026

Keywords: antidepressant activity, depression, serotonin, neuroplasticity, bdnf, glutamate, behavioural models, oxidative stress, glutamate; oxidative stress; neuroinflammation

Abstract

Depression is a prevalent and complex neuropsychiatric disorder characterized by persistent low mood, anhedonia, cognitive impairment, sleep disturbances, and alterations in motivation and behavior. Although conventional antidepressant drugs primarily target monoaminergic neurotransmission, increasing evidence indicates that the biological basis of antidepressant activity extends beyond serotonin, norepinephrine, and dopamine signaling. Antidepressant activity involves a network of interconnected mechanisms, including regulation of neuroplasticity, brain-derived neurotrophic factor signaling, hypothalamic–pituitary–adrenal axis activity, inflammation, oxidative stress, glutamatergic neurotransmission, and neuronal connectivity. Experimental evaluation of antidepressant activity commonly employs behavioral models such as the forced swim test, tail suspension test, sucrose preference test, and learned helplessness paradigm. These models provide useful information about behavioral responses to candidate compounds, although their interpretation requires consideration of their limitations and relevance to human depression. Conventional pharmacological agents, including selective serotonin reuptake inhibitors, serotonin–norepinephrine reuptake inhibitors, tricyclic antidepressants, and atypical antidepressants, have demonstrated antidepressant effects through different neurochemical pathways.

Introduction

Depression is one of the most important mental health disorders worldwide and can substantially affect emotional well-being, cognition, social functioning, and quality of life. It is a multifactorial condition influenced by genetic, environmental, psychological, neurochemical, endocrine, and inflammatory factors. The complexity of depression has encouraged

researchers to investigate multiple biological pathways rather than relying on a single neurotransmitter hypothesis.

Antidepressant activity refers to the ability of a substance or therapeutic intervention to reduce depression-related symptoms or produce behavioral and neurobiological changes associated with antidepressant effects. In experimental research, antidepressant activity is often evaluated by observing changes in behavioral responses following administration of a test compound

Neurobiological Basis of Antidepressant Activity

Monoaminergic neurotransmission

Serotonin, norepinephrine, and dopamine play important roles in mood regulation, motivation, reward processing, attention, and emotional responses. Selective serotonin reuptake inhibitors (SSRIs) increase serotonergic signaling primarily by inhibiting the serotonin transporter. Serotonin–norepinephrine reuptake inhibitors (SNRIs) enhance both serotonergic and noradrenergic neurotransmission. Tricyclic antidepressants also influence these neurotransmitter systems but generally have broader receptor activity. Changes in monoaminergic signaling may subsequently influence intracellular signaling pathways and neuronal adaptation, which may contribute to the delayed clinical effects of conventional antidepressants.

Neuroplasticity and BDNF

Neuroplasticity describes the capacity of the nervous system to modify neuronal structure and function in response to internal and external stimuli. Chronic stress and depression have been associated with

• [Journal of Neuroscience and Neurodevelopmental Disorders \(JNDD\)](#)

terations in neuronal connectivity and plasticity in several brain regions.

Brain-derived neurotrophic factor (BDNF) is an important mediator of neuronal survival, synaptic function, and plasticity. Antidepressant treatments may influence BDNF-related signaling and downstream pathways involved in neuronal adaptation. The interaction between antidepressant treatment and neuroplasticity has therefore become an important area of research.

Experimental Evaluation of Antidepressant Activity

Animal and laboratory models are frequently used during early-stage antidepressant research. These models are useful for identifying behavioral and biological effects of candidate compounds but cannot reproduce the full complexity of human depression.

Forced Swim Test

The forced swim test is widely used to assess behavioral responses to antidepressant treatments in rodents. Animals are placed in a container containing water, and changes in behavioral strategies are recorded. A reduction in immobility following administration of a test compound has traditionally been interpreted as an antidepressant-like behavioral effect. However, immobility can reflect multiple processes, and the test should not be considered a direct model of human depression.

Tail Suspension Test

The tail suspension test is another commonly used behavioral paradigm. Rodents are suspended by their tails, and periods of immobility are measured. Certain antidepressant treatments reduce immobility under standardized experimental conditions. Like the forced swim test, this assay is primarily a screening tool rather than a complete model of depressive illness.

Sucrose Preference Test

Anhedonia, or reduced ability to experience pleasure, is an important feature of depression. The sucrose preference test evaluates changes in the preference for a sweet solution compared with water. Reduced sucrose preference can be used as an indicator of anhedonia-like behavior in experimental models. Restoration of sucrose preference following treatment may provide evidence of an antidepressant-like effect.

Conclusion

Antidepressant activity is a complex biological phenomenon involving multiple interconnected

pathways rather than a single neurotransmitter system. Although monoaminergic mechanisms remain important, increasing evidence highlights the roles of neuroplasticity, BDNF signaling, glutamatergic neurotransmission, HPA-axis regulation, inflammation, and oxidative stress.

Experimental models such as the forced swim test, tail suspension test, sucrose preference test, and chronic stress paradigms are valuable tools for preliminary investigation, but their limitations must be recognized. Continued research into diverse molecular pathways and improved experimental models may contribute to the development of more effective and individualized antidepressant therapies.

References

1. Lee, J.H.; Lee, D.U.; Jeong, C.S. Gardenia jasminoides Ellis ethanol extract and its constituents reduce the risks of gastritis and reverse gastric lesions in rats. *Food Chem. Toxicol.* 2009, 47, 1127–1131.
2. Khushboo; Siddiqi, N.J.; de Lourdes Pereira, M.; Sharma, B. Neuroanatomical, Biochemical, and Functional Modifications in Brain Induced by Treatment with Antidepressants. *Mol. Neurobiol.* 2022, 59, 3564–3584.
3. Macrea, C.; Ilias, T.; Fratila, O.; Roxana, B.; Hocopan, C. Probiotics and the microbiota-gut-brain axis: Focus on patients with depression. A review of current research. *Biomed Pap. Med. Fac. Univ. Palacky Olomouc Czech Repub.* 2023, 167, 219–224.
4. Mayer, E.A.; Tillisch, K.; Gupta, A. Gut/brain axis and the microbiota. *J. Clin. Invest.* 2015, 125, 926–938.
5. Dong, W.R.; Liu, H.Y.; Ding, Y.G.; Wang, H.P.; Liu, S.M. Interpretation of a TCM theory about clearing and discharging the triple burner damp-heat of Gardenia jasminoides Ellis. *China J. Trad. Chin. Med. Pharm.* 2011, 26, 33–35.

- [Journal of Neuroscience and Neurodevelopmental Disorders \(JNDD\)](#)



This work is licensed under Creative Commons Attribution 4.0 License
DOI:10/JNDD/2026/023

Your next submission with

Olites Publishers will reach you the below assets

- We follow principles of publication led by the Committee on Publication Ethics (COPE).
- Double blinded peer review process which is just as well as constructive.
- Permanent archiving of your article on our website
- Quality Editorial service
- Manuscript accessibility in different formats (PDF, Full Text)
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

- Learn more: [Journal of Neuroscience and Neurodevelopmental Disorders](#)
[Olites Publishers \(olitespublishing.org\)](#)